

All Databases

PubMed

Nucleotide

Protein

Genome

Structure

OMIM

PMC

Journal

Search for

Limits

Preview/Index

History

Clipboard

Details

Display

Summary

Show:

Sort

Send to

Text

About Entrez

Text Version

Items 1 - 46 of 46

One p

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed

Services

Journals

Database

MeSH Database

Single Citation

Matcher

Batch Citation

Matcher

Clinical Queries

LinkOut

My NCBI

(Cubby)

Related

Resources

Order

Documents

NLM Catalog

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

☐ 1: [Dong M, Pinon DI, Miller LJ.](#)

Related Articles, I



Insights into the structure and molecular basis of ligand docking to the G protein-coupled secretin receptor using charge-modified amino terminal agonist probes.

Mol Endocrinol. 2005 Feb 24; [Epub ahead of print]

PMID: 15731172 [PubMed - as supplied by publisher]

☐ 2: [Lee SM, Yung WH, Chen L, Chow BK.](#)

Related Articles, I



Expression and spatial distribution of secretin and secretin receptor in human cerebellum.

Neuroreport. 2005 Feb 28;16(3):219-22.

PMID: 15706223 [PubMed - in process]

☐ 3: [Langer I, Gregoire F, Nachtergaeel I, De Neef P, Vertongen P, Robberecht P.](#)

Related Articles, I



Hexanoylation of a VPAC2 receptor-preferring ligand markedly increased its selectivity and potency.

Peptides. 2004 Feb;25(2):275-8.

PMID: 15063009 [PubMed - indexed for MEDLINE]

☐ 4: [Mustafa T, Klonisch T, Hombach-Klonisch S, Kehlen A, Schmutzler C, Koehrle J, Gimm O, Dralle H, Hoang-Vu C.](#)

Related Articles, I



Expression of CD97 and CD55 in human medullary thyroid carcinomas.

Int J Oncol. 2004 Feb;24(2):285-94.

PMID: 14719104 [PubMed - indexed for MEDLINE]

☐ 5: [Dong M, Li Z, Pinon DI, Lybrand TP, Miller LJ.](#)

Related Articles, I



Spatial approximation between the amino terminus of a

peptide agonist and the top of the sixth transmembrane segment of the secretin receptor.

J Biol Chem. 2004 Jan 23;279(4):2894-903. Epub 2003 Oct 30.
PMID: 14593094 [PubMed - indexed for MEDLINE]

- ▢ 6: Dong M, Li Z, Zang M, Pinon DI, Lybrand TP, Miller LJ. Related Articles, I



Spatial approximation between two residues in the mid-region of secretin and the amino terminus of its receptor. Incorporation of seven sets of such constraints into a three dimensional model of the agonist-bound secretin receptor. J Biol Chem. 2003 Nov 28;278(48):48300-12. Epub 2003 Sep 18
PMID: 14500709 [PubMed - indexed for MEDLINE]

- ▢ 7: Taylor WR, Munro RE, Petersen K, Bywater RP. Related Articles, I



Ab initio modelling of the N-terminal domain of the secretin receptors.

Comput Biol Chem. 2003 May;27(2):103-14.
PMID: 12821307 [PubMed - indexed for MEDLINE]

- ▢ 8: Zang M, Dong M, Pinon DI, Ding XQ, Hadac EM, Li Z, Lybrand TP, Miller LJ. Related Articles, I



Spatial approximation between a photolabile residue in position 13 of secretin and the amino terminus of the secretin receptor.

Mol Pharmacol. 2003 May;63(5):993-1001. Erratum in: Mol Pharmacol. 2003 Jun;64(1):193.
PMID: 12695527 [PubMed - indexed for MEDLINE]

- ▢ 9: Mayo KE, Miller LJ, Bataille D, Dalle S, Goke B, Thorens B, Drucker DJ. Related Articles, I



International Union of Pharmacology. XXXV. The glucagon receptor family.

Pharmacol Rev. 2003 Mar;55(1):167-94. Review.
PMID: 12615957 [PubMed - indexed for MEDLINE]







- ▢ 10: Dong M, Miller LJ. Related Articles, I



Molecular pharmacology of the secretin receptor.


Receptors Channels. 2002;8(3-4):189-200. Review.
PMID: 12529936 [PubMed - indexed for MEDLINE]

- ▢ 11: Cunha SR, Mayo KE. Related Articles, I


-  Ghrelin and growth hormone (GH) secretagogues potentiate GH-releasing hormone (GHRH)-induced cyclic adenosine 3',5'-monophosphate production in cells expressing transfected GHRH and GH secretagogue receptors.
Endocrinology. 2002 Dec;143(12):4570-82.
PMID: 12446584 [PubMed - indexed for MEDLINE]
- ☐ **12:** Joost P, Methner A. Related Articles, 1
-  Phylogenetic analysis of 277 human G-protein-coupled receptors as a tool for the prediction of orphan receptor ligands.
Genome Biol. 2002 Oct 17;3(11):RESEARCH0063. Epub 2002 Oct 17.
PMID: 12429062 [PubMed - indexed for MEDLINE]
- ☐ **13:** Dong M, Pinon DI, Miller LJ. Related Articles, 1
-  Development of a biologically active secretin analogue incorporating a radioiodinatable photolabile p-(4-hydroxybenzoyl)phenylalanine in position 10.
Regul Pept. 2002 Nov 15;109(1-3):181-7.
PMID: 12409231 [PubMed - indexed for MEDLINE]
- ☐ **14:** Unson CG, Wu CR, Jiang Y, Yoo B, Cheung C, Sakmar TP, Merrifield RB. Related Articles, 1
-  Roles of specific extracellular domains of the glucagon receptor in ligand binding and signaling.
Biochemistry. 2002 Oct 1;41(39):11795-803.
PMID: 12269822 [PubMed - indexed for MEDLINE]
- ☐ **15:** Waelbroeck M, Perret J, Vertongen P, Van Craenenbroeck M, Robberecht P. Related Articles, 1
-  Identification of secretin, vasoactive intestinal peptide and glucagon binding sites: from chimaeric receptors to point mutations.
Biochem Soc Trans. 2002 Aug;30(4):437-41. Review.
PMID: 12196110 [PubMed - indexed for MEDLINE]
- ☐ **16:** Du K, Couvineau A, Rouyer-Fessard C, Nicole P, Laburthe M. Related Articles, 1
-  Human VPAC1 receptor selectivity filter. Identification of a critical domain for restricting secretin binding.

J Biol Chem. 2002 Oct 4;277(40):37016-22. Epub 2002 Jul 19.
PMID: 12133828 [PubMed - indexed for MEDLINE]


- ▣ **17:** Perret J, Van Craenenbroeck M, Langer I, Vertongen P, Gregoire F, Robberecht P, Waelbroeck M. Related Articles, 1

 Mutational analysis of the glucagon receptor: similarities with the vasoactive intestinal peptide (VIP)/pituitary adenylate cyclase-activating peptide (PACAP)/secretin receptors for recognition of the ligand's third residue. Biochem J. 2002 Mar 1;362(Pt 2):389-94. PMID: 11853547 [PubMed - indexed for MEDLINE]


- ▣ **18:** Chan KY, Pang RT, Chow BK. Related Articles, 1

 Functional segregation of the highly conserved basic motifs within the third endolobe of the human secretin receptor. Endocrinology. 2001 Sep;142(9):3926-34. PMID: 11517171 [PubMed - indexed for MEDLINE]


- ▣ **19:** West AP Jr, Llamas LL, Snow PM, Benzer S, Bjorkman PJ. Related Articles, 1

 Crystal structure of the ectodomain of Methuselah, a Drosophila G protein-coupled receptor associated with extended lifespan. Proc Natl Acad Sci U S A. 2001 Mar 27;98(7):3744-9. Epub 2001 Mar 13. PMID: 11274391 [PubMed - indexed for MEDLINE]

- ▣ **20:** Dong M, Wang Y, Miller LJ. Related Articles, 1

 Dual contacts between peptide agonist ligands and the secretin receptor directly established by photoaffinity labeling. Ann N Y Acad Sci. 2000;921:381-6. PMID: 11193860 [PubMed - indexed for MEDLINE]

- ▣ **21:** Park CG, Ganguli SC, Pinon DI, Hadac EM, Miller LJ. Related Articles, 1

 Cross-chimeric analysis of selectivity of secretin and VPAC(1) receptor activation. J Pharmacol Exp Ther. 2000 Nov;295(2):682-8. PMID: 11046106 [PubMed - indexed for MEDLINE]

- ▣ **22:** Dong M, Asmann YW, Zang M, Pinon DI, Miller LJ. Related Articles, 1



Identification of two pairs of spatially approximated residues within the carboxyl terminus of secretin and its receptor.

J Biol Chem. 2000 Aug 25;275(34):26032-9.

PMID: 10859300 [PubMed - indexed for MEDLINE]

- ▣ **23:** Di Paolo E, Vilardaga JP, Petry H, Moguilevsky N, Bollen A, Robberecht P, Waelbroeck M. Related Articles, 1



Role of charged amino acids conserved in the vasoactive intestinal polypeptide/secretin family of receptors on the secretin receptor functionality.

Peptides. 1999;20(10):1187-93.

PMID: 10573290 [PubMed - indexed for MEDLINE]

- ▣ **24:** Di Paolo E, De Neef P, Moguilevsky N, Petry H, Cnudde J, Bollen A, Waelbroeck M, Robberecht P. Related Articles, 1



Properties of a recombinant human secretin receptor: a comparison with the rat and rabbit receptors.

Pancreas. 1999 Jul;19(1):51-5.

PMID: 10416692 [PubMed - indexed for MEDLINE]

- ▣ **25:** Di Paolo E, Petry H, Moguilevsky N, Bollen A, De Neef P, Waelbroeck M, Robberecht P. Related Articles, 1



Mutations of aromatic residues in the first transmembrane helix impair signalling by the secretin receptor.

Receptors Channels. 1999;6(4):309-15.

PMID: 10412723 [PubMed - indexed for MEDLINE]

- ▣ **26:** Frimurer TM, Bywater RP. Related Articles, 1



Structure of the integral membrane domain of the GLP1 receptor.

Proteins. 1999 Jun 1;35(4):375-86.

PMID: 10382665 [PubMed - indexed for MEDLINE]

- ▣ **27:** Dong M, Wang Y, Hadac EM, Pinon DI, Holicky E, Miller LJ. Related Articles, 1



Identification of an interaction between residue 6 of the natural peptide ligand and a distinct residue within the

amino-terminal tail of the secretin receptor.

J Biol Chem. 1999 Jul 2;274(27):19161-7.

PMID: 10383421 [PubMed - indexed for MEDLINE]

▢ **28:** Olde B, Sabirsh A, Owman C. Related Articles, 1



Molecular mapping of epitopes involved in ligand activation of the human receptor for the neuropeptide, based on hybrids with the human secretin receptor.

J Mol Neurosci. 1998 Oct;11(2):127-34.

PMID: 10096039 [PubMed - indexed for MEDLINE]

▢ **29:** Ng SS, Pang RT, Chow BK, Cheng CH. Related Articles, 1



Real-time evaluation of human secretin receptor activity using cytosensor microphysiometry.

J Cell Biochem. 1999 Mar 15;72(4):517-27.

PMID: 10022611 [PubMed - indexed for MEDLINE]

▢ **30:** Dong M, Wang Y, Pinon DI, Hadac EM, Miller LJ. Related Articles, 1



Demonstration of a direct interaction between residue 2 of the carboxyl-terminal half of secretin and the amino-terminal tail of the secretin receptor using photoaffinity labeling.

J Biol Chem. 1999 Jan 8;274(2):903-9.

PMID: 9873030 [PubMed - indexed for MEDLINE]

▢ **31:** Ganguli SC, Park CG, Holtmann MH, Hadac EM, Kenakin TP, Miller LJ. Related Articles, 1



Protean effects of a natural peptide agonist of the G protein-coupled secretin receptor demonstrated by receptor mutagenesis.

J Pharmacol Exp Ther. 1998 Aug;286(2):593-8.

PMID: 9694908 [PubMed - indexed for MEDLINE]

▢ **32:** DeAlmeida VI, Mayo KE. Related Articles, 1




Identification of binding domains of the growth hormone releasing hormone receptor by analysis of mutant and chimeric receptor proteins.

Mol Endocrinol. 1998 May;12(5):750-65.


PMID: 9605937 [PubMed - indexed for MEDLINE]


▢ **33:** Di Paolo E, De Neef P, Moguilevsky N. Related Articles, 1


Petry H, Bollen A, Waelbroeck M, Robberecht P.

-  Contribution of the second transmembrane helix of the secretin receptor to the positioning of secretin.
FEBS Lett. 1998 Mar 13;424(3):207-10.
PMID: 9539152 [PubMed - indexed for MEDLINE]


 **34:** Chow BK. Related Articles, 1

-  Functional antagonism of the human secretin receptor by recombinant protein encoding the N-terminal ectodomain of the receptor.
Recept Signal Transduct. 1997;7(3):143-50.
PMID: 9440501 [PubMed - indexed for MEDLINE]


 **35:** Gourlet P, Vandermeers A, Vandermeers-Piret MC, De Neef P, Waelbroeck M, Robberecht P. Related Articles, 1


-  Effect of introduction of an arginine16 in VIP, PACAP secretin on ligand affinity for the receptors.
Biochim Biophys Acta. 1996 Dec 12;1314(3):267-73.
PMID: 8982281 [PubMed - indexed for MEDLINE]


 **36:** van Lier RA, Eichler W, Hamann J. Related Articles, 1

-  Seven-span transmembrane molecules: novel receptors involved in leukocyte adhesion.
Immunol Lett. 1996 Dec;54(2-3):185-7. Review.
PMID: 9052875 [PubMed - indexed for MEDLINE]

 **37:** Mullaney I, Carr IC, Milligan G. Related Articles, 1

-  Overexpression of G(s)alpha in NG108-15, neuroblastomaXglioma cells: effects on receptor regulation of the stimulatory adenylyl cyclase cascade.
FEBS Lett. 1996 Nov 18;397(2-3):325-30.
PMID: 8955373 [PubMed - indexed for MEDLINE]

 **38:** Holtmann MH, Hadac EM, Ulrich CD, Miller LJ. Related Articles, 1

-  Molecular basis and species specificity of high affinity binding of vasoactive intestinal polypeptide by the rat secretin receptor.
J Pharmacol Exp Ther. 1996 Nov;279(2):555-60.
PMID: 8930157 [PubMed - indexed for MEDLINE]

- ▢ **39:** Hamann J, Vogel B, van Schijndel GM, van Lier RA. Related Articles, 1



The seven-span transmembrane receptor CD97 has a cellular ligand (CD55, DAF).

J Exp Med. 1996 Sep 1;184(3):1185-9.

PMID: 9064337 [PubMed - indexed for MEDLINE]

- ▢ **40:** Holtmann MH, Ganguli S, Hadac EM, Dolu V, Miller LJ. Related Articles, 1



Multiple extracellular loop domains contribute critical determinants for agonist binding and activation of the secretin receptor.

J Biol Chem. 1996 Jun 21;271(25):14944-9.

PMID: 8663161 [PubMed - indexed for MEDLINE]

- ▢ **41:** Turner PR, Bambino T, Nissenson RA. Related Articles, 1



A putative selectivity filter in the G-protein-coupled receptors for parathyroid hormone and secretion.

J Biol Chem. 1996 Apr 19;271(16):9205-8.

PMID: 8621578 [PubMed - indexed for MEDLINE]

- ▢ **42:** Vilardaga JP, di Paolo E, de Neef P, Waelbroeck M, Bollen A, Robberecht P. Related Articles, 1



Lysine 173 residue within the first exoloop of rat secretin receptor is involved in carboxylate moiety recognition (Asp 3 in secretin).

Biochem Biophys Res Commun. 1996 Jan 26;218(3):842-6.

PMID: 8579602 [PubMed - indexed for MEDLINE]

- ▢ **43:** Ozcelebi F, Holtmann MH, Rentsch RU, Rao R, Miller LJ. Related Articles, 1



Agonist-stimulated phosphorylation of the carboxyl-terminal tail of the secretin receptor.

Mol Pharmacol. 1995 Nov;48(5):818-24.

PMID: 7476911 [PubMed - indexed for MEDLINE]

- ▢ **44:** Vilardaga JP, De Neef P, Di Paolo E, Bollen A, Waelbroeck M, Robberecht P. Related Articles, 1



Properties of chimeric secretin and VIP receptor protein indicate the importance of the N-terminal domain for ligand discrimination.

Biochem Biophys Res Commun. 1995 Jun 26;211(3):885-91.

PMID: 7598719 [PubMed - indexed for MEDLINE]

- ☐ **45:** [Usdin TB, Mezey E, Button DC, Brownstein MJ, Bonner TI.](#) Related Articles, 1



Gastric inhibitory polypeptide receptor, a member of the secretin-vasoactive intestinal peptide receptor family, is widely distributed in peripheral organs and the brain.

Endocrinology. 1993 Dec;133(6):2861-70.

PMID: 8243312 [PubMed - indexed for MEDLINE]

- ☐ **46:** [Roth BL, Beinfeld MC, Howlett AC.](#) Related Articles, 1



Secretin receptors on neuroblastoma cell membranes: characterization of 125I-labeled secretin binding and association with adenylate cyclase.

J Neurochem. 1984 Apr;42(4):1145-52.

PMID: 6321661 [PubMed - indexed for MEDLINE]

Display Show: Sort Send to

[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

Mar 14 2005 07:08:36